

Diarrhoea and vomiting

Not nice for you or your baby

Sickness and diarrhoea bugs are caught easily and are often passed on in places where there are lots of children. They are generally caused by viruses and those affected (adults and children) recover within 48 hours. It is important to drink plenty of fluid to prevent becoming dehydrated.

Feeling sick and suddenly being sick are normally the first signs. Diarrhoea can follow afterwards. If your child is not vomiting frequently, is reasonably comfortable in between and you are able to give them frequent small amounts of water, they are less likely to become dehydrated and probably don't need to see a doctor. **Speak to your GP if they are unwell for longer than 24 hours or sooner if they are newborn or if you notice signs of dehydration.**

Continue breastfeeding or other milk feeds - do so even more frequently. Offer older children plenty of water, or an ice-lolly for them to suck. If they want to eat, give them plain foods like pasta or boiled rice (nothing too rich or salty).

Keep them away from others, especially children and older people, who may pick up infection. Be extra careful with everyone's handwashing.

Pharmacist says

There are lots of ways you can care for your child at home. Things to try are:

- Give them regular drinks - try small amounts of boiled cooled water if bottle fed.
- If breastfeeding, carry on as usual. If not, continue with other milk feeds.
- Being extra careful with hand hygiene (use soap and water and dry hands well with a clean towel).
- Rehydrating solutions come in pre-measured sachets to mix with boiled cooled water. It helps with dehydration.

If your child is unwell for more than 24 hours speak to your GP. If your baby is newborn or very unwell contact your GP straight away.

Signs of dehydration

- Less wet nappies.
- More sleepy than usual.
- Dry mouth.
- Sunken fontanelle (the soft spot on the top of the head is more dipped in than usual).

Try a rehydrating solution from your pharmacist.

1

My baby has diarrhoea and is being sick.

2

Have you given them lots of fluids? This will help prevent them becoming dehydrated if it is a tummy bug. Speak to your pharmacist and ask about a rehydrating solution.

3

Speak to your GP if symptoms show no sign of improvement after 24 hours or straight away if they are newborn.